

The risk that sits between specialties

Medicine is largely organised around organs. Chronic obstructive pulmonary disease (COPD) belongs to respiratory medicine; myocardial infarction, heart failure and arrhythmia belong to cardiology.

However, the patient does not experience these diseases in isolation.
Neither respiratory nor cardiology fully owns this important interface. As a result, the cardiovascular consequences of COPD exacerbations can fall between traditional specialty silos. This matters because cardiovascular disease is one of the leading causes of illness and death in COPD, particularly in patients with mild-to-moderate disease.

General practitioners are uniquely placed to see what an organ-specific model can miss.



An exacerbation is a systemic event

A COPD exacerbation is often viewed as a temporary increase in breathlessness, cough or sputum. Treat the infection or airway inflammation, wait for the symptoms to settle, and the episode appears to be over.

Biologically, considerably more is happening.

An exacerbation may produce systemic inflammation, increased sympathetic activity, hypoxaemia, endothelial dysfunction and activation of platelets and coagulation pathways. Infection, tachycardia and increased respiratory effort place further physiological stress on an already vulnerable cardiovascular system. This creates a period in which heart failure, arrhythmia, myocardial infarction and stroke become more likely. Heart failure and arrhythmia appear to be particularly important—not just the traditional concern about acute coronary events.

The danger continues after the lungs improve

The cardiovascular risk is greatest during the first days and weeks after an exacerbation, but it does not immediately return to baseline when the respiratory symptoms settle.

In one large study of patients with COPD and established or increased cardiovascular risk, the likelihood of a major cardiovascular event was almost four times higher during the first 30 days after an exacerbation. After an exacerbation requiring hospitalisation, it was nearly ten times higher.

Severe exacerbations carry the greatest risk, but this is not only a hospital phenomenon. Exacerbations treated in general practice with antibiotics or

corticosteroids are also followed by a measurable increase in myocardial infarction and stroke.

Not every subsequent cardiovascular event is necessarily caused by the exacerbation. Shared risk factors, underlying disease and diagnostic overlap all contribute. Nevertheless, the consistent timing and severity gradient across large studies make the association difficult to dismiss.

One exacerbation predicts the next

An exacerbation is not simply an isolated event. A previous treated exacerbation is the strongest predictor of a future exacerbation, even after accounting for the severity of airflow obstruction.

This may partly reflect an exacerbation-prone phenotype rather than one event directly causing another. Clinically, however, the effect is the same: patients who exacerbate are more likely to experience another episode—and therefore another period of systemic and cardiovascular vulnerability.

The familiar respiratory cycle is therefore also a cardiovascular cycle:



The GP advantage

The practical insight is not another assessment checklist. It is a change in how we interpret an exacerbation.

When a patient with COPD experiences an exacerbation, the event should not be considered closed simply because the cough and breathlessness have improved. It identifies a patient at increased risk of another exacerbation and a period of increased cardiovascular vulnerability.

The specialist view divides disease by organs. The GP sees the whole patient. At this interface, that broader perspective may be exactly what is needed.

The exacerbation may be over—but the patient's respiratory and cardiovascular risk has not necessarily returned to baseline.

WRS FOR GP

Evidence. Insight. Precision.

WRS For GP, Evidence Insight Precision, Western Respiratory Sleep, Joondalup, WA 6027, Australia

Unsubscribe Manage preferences